



PRESS-RELEASE

Max Planck Institute of Quantum Optics and Munich-Centre for Advanced Photonics




Garching, 27 October, 2015

Film production in 4D with ultrashort electron pulses

Physicists of the Ludwig-Maximilians-Universität Munich and the Max Planck Institute of Quantum Optics shorten electron pulses down to 30 femtoseconds duration. This enables them to gain detailed insight into atomic motions in molecules.

Electrons are odd particles: they have both wave and particle properties. Electron microscopy has been taking advantage of this phenomenon for roughly a century now and grants us a direct insight into the fundamental components of matter: molecules and atoms. For a long time, still images were provided, but for some years now scientists are making tremendous progress in short-pulse technology. They create beams of electron pulses, which can, due to their extremely short flashing, provide us with very sharp images of moving atoms and electrons. Nevertheless, some of the fastest processes still remained blurred. A team of the Laboratory for Attosecond Physics (LAP) from the Ludwig-Maximilians-Universität (LMU) and the Max Planck Institute of Quantum Optics (MPQ) has now managed to shorten electron pulses down to 28 femtoseconds in duration. One femtosecond is a millionth of a billionth of a second (10^{-15} s). Such shutter speeds enable us to directly observe the truly fundamental motions of atoms and molecules in solids, similar to stroboscopy.



*Figure: If ultrashort electron pulses hit a biomolecular crystal, they are diffracted from it. As a result, one obtains a characteristic diffraction image of the atomic structure. **Graphic: Alexander Gliserin***

Those who want to explore the microcosm and its dynamics need a high-speed camera for atoms. In order to sharply capture motions of such particles during a reaction, one needs to work with “shutter speeds” in the range of femtoseconds, since this is the speed of reactions in molecules and solids. Commonly, femtosecond-short shutter speeds are provided by short-

pulse laser technology, but laser light is not able to spatially resolve atoms. Scientists from the Laboratory for Attosecond Physics at LMU and MPQ have now succeeded in producing ultrashort electron pulses with a duration of only 28 femtoseconds. This is six times shorter than ever before. The length of the matter wave is only about eight picometers; one picometer is a trillionth of a meter (10^{-12} m). Due to this short wavelength, it is possible to visualize even single atoms in diffraction experiments. If such electrons meet a molecule or atom, they are diffracted into specific directions due to their short wavelength. This way they generate an interference pattern at the detector from

Max Planck Institute of Quantum Optics
Press & Public Relations
Dr. Olivia Meyer-Streng
Phone: +49-89-32905-213
E-mail: olivia.meyer-streng@mpq.mpg.de
Hans-Kopfermann-Str. 1, D-85748 Garching

Munich-Centre for Advanced Photonics
Public Outreach
Thorsten Naeser
Phone: +49-89-32905-124
E-mail: thorsten.naeser@mpq.mpg.de

which an atomic 3D-structure of the examined substance is reconstructed. If the pulses are short enough, a sharp snapshot of the movement is the result.

To test the new technique, the physicists applied their ultrashort electron pulses to a biomolecule in a diffraction experiment. It is planned to use those electron beams for pump-probe experiments: an optical laser pulse is sent to the sample, initiating a response. Shortly afterwards the electron pulses produce a diffraction image of the structure at a sharp instant in time. A large amount of such snapshots at varying delay times between the initiating laser pulses and the electron pulses then results in a film showing the atomic motion within the substance. Thanks to the sub-atomic wavelength of the electrons, one therefore obtains a spatial image as well as the dynamics. Altogether this results in a four-dimensional impression of molecules and their atomic motions during a reaction.

„With our ultrashort electron pulses, we are now able to gain a much more detailed insight into processes happening within solids and molecules than before“, Dr. Peter Baum says. „We are now able to record the fastest known atomic motions in four dimensions, namely in space and time“. Now the physicists aim to further reduce the duration of their electron pulses. The shorter the shutter speed becomes, the faster the motions which can be recorded. The aim of the scientists is to eventually observe even the much faster motions of electrons in light-driven processes.

Thorsten Naeser

Original publication:

A. Gliserin, M. Walbran, F. Krausz, P. Baum

Sub-phonon-period compression of electron pulses for atomic diffraction

Nature Communications, 27 October 2015, doi: 10.1038/ncomms9723

For further information, contact:

Dr. Peter Baum

Max Planck Institute of Quantum Optics
Ludwig-Maximilians-Universität Munich
Am Coulombwall 1, 85748 Garching, Germany
Phone: +49 (0)89 289 - 14102
E-mail: peter.baum@lmu.de

Prof. Ferenc Krausz

Chair of Experimental Physics,
Ludwig-Maximilians-Universität Munich
Laboratory for Attosecond Physics
Director at Max Planck Institute of Quantum Optics
Hans-Kopfermann-Str. 1, 85748 Garching, Germany
Phone: +49 (0)89 32 905 - 600 / Fax: - 649
E-mail: ferenc.krausz@mpq.mpg.de
www.attoworld.de